

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representation of
The original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 721 322 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:
18.10.2000 Bulletin 2000/42

(21) Application number: 92905356.9

(22) Date of filing: 14.01.1992

(51) Int. Cl.⁷: **A61K 7/15**

(86) International application number:
PCT/US92/00409

(87) International publication number:
WO 92/12700 (06.08.1992 Gazette 1992/21)

(54) **SHAVING COMPOSITIONS**

RASIERPRÄPARATE

COMPOSITIONS DE RASAGE

(84) Designated Contracting States:
AT BE CH DE DK ES FR GB GR IT LI LU MC NL
SE

(30) Priority: 25.01.1991 US 645938

(43) Date of publication of application:
17.07.1996 Bulletin 1996/29

(73) Proprietor: **THE GILLETTE COMPANY**
Boston, Massachusetts 02199 (US)

(72) Inventors:

- BARNET, Alfred, G.
Hingham, MA 02043 (US)
- MEZIKOFSKY, Merrill, R.
Wakefield, MA 01880 (US)

(74) Representative:
Baillie, Iain Cameron
Ladas & Parry,
Dachauerstrasse 37
80335 München (DE)

(56) References cited:
US-A- 3 912 666 US-A- 4 528 111
US-A- 4 765 975

Remarks:

The file contains technical information submitted
after the application was filed and not included in
this specification

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description

[0001] This invention relates to improved shaving preparations of the type dispensed in the form of a gel containing a volatile component that causes the gel to turn into a foam by itself when spread on the skin in preparation for wet shaving, i.e., shaving with a razor blade.

[0002] The class of shaving preparation referred to herein as a "self-foaming gel" is a lather-producing composition which is capable of being dispensed as a gel that is substantially free from foaming, but which when spread over the skin and beard produces a foam generated by the action of a volatile self-foaming agent. Such compositions generally take the form of an oil-in-water emulsion in which the self-foaming agent, generally a low-boiling aliphatic hydrocarbon, solubilizes the oil phase, and the water phase comprises a water-soluble soap component so selected that the resulting packaged composition is in the form of a stable gel under ambient conditions.

[0003] Such products are typically packaged in an aerosol container with a barrier such as a piston or collapsible bag to separate the self-foaming gel from the propellant required for expulsion and thereby ensure that a homogeneous gel is dispensed, substantially free of bubbles, which produce a self-generated lather of uniform consistency and density throughout the life of the product. Other known packaging forms for self-foaming gels include conventional, i.e., non-barrier, aerosol containers and collapsible tubes, although these less expensive packages have not always been entirely satisfactory due to formulation compromises attendant with their use.

[0004] For further description of parameters involved in the formulation of such products, reference may be made, for example, to U.S. Patents 3,541,581 and 4,528,111; typical manufacturing processes and equipment are disclosed in, for example, U.S. Patents 4,405,489 and 4,651,503. Other references include US-A-4 765 975 which discloses the use of fluorosurfactants in hair conditioning compositions; US-A-3 912 666 which discloses the use of self-foaming agents in aerosol cosmetic applications; and Lexikon der Hilfsstoffe..., H.P. Fiedler, 1989, Seite 996 which discloses certain characteristics of Polysynlane,

[0005] Self-foaming shaving gels are, as noted, well known in the art and have been commercially available for over twenty years, during which time they have achieved a substantial share of the total market for shave preparations. Users tend to exhibit considerable loyalty to this form of shaving preparation and associate its use with obtaining a close, comfortable shave. Nevertheless, further improvements in performance would be desirable.

[0006] We have now unexpectedly found a significant improvement in certain key performance attributes of shaving gels comprising water, a water-soluble soap component, and a self-foaming agent, which additionally include from about 0.0005 to 0.5 percent by weight of a fluorosurfactant in combination with a hydrogenated polyisobutene. In preferred embodiments of the invention lubricity is enhanced by the inclusion of one or more water-soluble polymers, and the composition may also include such optional components as clarity aids, foam modifiers, humectants, perfumes, colorants, skin conditioners, and emollients.

[0007] The major essential component of the shaving gel is about 55-85%, preferably about 65-80% by weight of water, which is preferably deionized or distilled water free of dissolved electrolytes such as salts and acids. The water component is essential to provide adequate solubility for the soap component and compatibility with the other essential ingredients and to provide a stable shaving gel having the desired foaming properties and capable of being readily rinsed from the skin.

[0008] The soap component of the composition, for the purposes hereof, may include a water-soluble salt of a fatty acid or any of certain, usually synthetic, wetting agents or detergents, or a combination thereof; where both types are used together, the combination is designated the soap component or, simply, the soap. All of these materials are well known in the art.

[0009] The water-soluble fatty acid salts may be preformed or formed in situ by reacting a basic material such as triethanolamine, sodium hydroxide, or potassium hydroxide with a higher fatty acid such as stearic, palmitic, myristic, oleic, coconut oil fatty acids, and mixtures thereof. The nature of the soap or detergent used is not critical but, in combination with the other components of the composition and their relative proportions, as is known in the art, has an effect on the properties of the shaving gel product of the invention and the lather produced. Particularly preferred are triethanolamine soaps of stearic and palmitic acids, though many and various animal and vegetable oil soaps and blends thereof are suitable for use in the practice of the invention.


[0010] The wetting agents that are useful as the soap component or as a portion of the soap component preferably are nonionic in character, and should be appreciably soluble in the aqueous components of the gel. Numerous such materials are well known in the art. As examples, mention may be made of such wetting agents as water-soluble polyoxyethylene ethers of alkyl-substituted phenols and water-soluble polyethoxylated derivatives of fatty alcohols. Particularly good results are obtained with Oleth-20, which when combined at a level of about 4-6% with suitable fatty acid soaps, results in the formation of a shaving gel composition of excellent properties without the need for any additional thickener or gelling agent. (Chemical names as used herein follow the system of nomenclature adopted by the Cosmetic, Toiletry and Fragrance Association, Inc.) The soap component constitutes about 5-30%, preferably about 12-25% by weight of the shaving gel.

[0011] Another essential ingredient of the self-foaming gel of the present invention is about 0.5-10% and preferably about 1-5% by weight of a volatile self-foaming agent which preferably is a hydrocarbon selected from the group consisting of n-pentane, isopentane, neopentane, n-butane, and isobutane, and mixtures thereof. A blend of isopentane and isobutane in a weight ratio of 3:1 is particularly desirable. Too much self-foaming agent can result in a product that is delivered as a foam, while too little results in greater effort and longer time to generate a foam on the skin.

[0012] Fluorosurfactants used in the present formulations are hydrophobic-lipophobic perfluorinated compounds which can be represented by the formula



wherein n is an integer from 4 to 18, R is selected from the group consisting of -X-, -Y-O-Z-, -Y-S-Z-,



-Y-CONH-Z-, -Y-SO₂NH-Z-, wherein X is (CH₂)_x, Y is (CH₂)_y and Z is (CH₂)_z, R₁ is an alkyl group containing from 1 to 4 carbon atoms, wherein x is an integer from 1 to 6, the terminal (CH₂)_y group is attached to the (C_nF_{2n+1}) portion of the molecule, the terminal (CH₂)_z group is attached to the Q portion of the molecule, Y is an integer from 0 to 3, z is an integer from 1 to 3, and Q is selected from the group of

-O-CH₂CH₂O)_xH, where x = 1 - 10;

-CO₂M, where M is an alkali metal (e.g., Na, K, Li);

-*N(CH₃)₃CH₂SO*

Commercially available surfactants which fall within the definition of this formula include Zonyl FSA[®], an anionic fluorochemical surfactant [R_fCH₂CH₂SCH₂CH₂CO₂Li] manufactured by E.I. DuPont de Nemours & Company; Zonyl FSK[®], an amphoteric fluorochemical surfactant [R_fCH₂CH(OCOCH₃)CH₂N⁺(CH₃)₂CH₂CO₂⁻] manufactured by E. I. DuPont de Nemours & Company; Zonyl FSN[®], a nonionic fluorochemical surfactant [R_fCH₂CH₂O(CH₂CH₂O)_xH], manufactured by E.I. DuPont de Nemours & Company where R_f = F(CF₂CF₂)₃₋₆; and Lodyne S-112B[®] - a blend of an anionic fluorochemical sodium sulfonate type and a nonionic fluorochemical synergist of the fluoroalkyl amide type manufactured by Ciba-Geigy. Cationic fluorosurfactants are not preferred, due to compatibility problems with any anionic soap component in the composition. Formulations of the invention contain a small but effective amount of fluorosurfactant, present at from about 0.0005 - 0.5 percent, preferably from about 0.001 - 0.1 percent and more preferably from about 0.002 - 0.01 weight percent of the composition as an active ingredient.

[0013] The hydrogenated polyisobutene component is represented by the formula [-CH₂-C(CH₃)₂]_n, where n (the number of repeating units) is about 6 to 24. An example of such material is Cosmetic Grade Panalane L-14E[®], which has an average molecular weight of about 320 dalton (i.e., n=6) from Amoco Chemical Company, Chicago, Illinois. Another suitable polyisobutene mixture is available from Nippon Oils under the name Polysynlane[®]. The hydrogenated polyisobutene constitutes about 0.5 - 5%, preferably about 1 - 4% by weight of the composition.

[0014] The self-foaming shaving gels of the present invention may, as indicated, form suitable gels without the inclusion of any thickener or gelling agent, depending particularly on the selection and concentration of the soap component of the composition. However, such "thickeners" as water-soluble polymers may be incorporated to impart enhanced lubricity to the compositions even without, in particular formulations, performing any thickening function. While water-soluble polymers as a class are useful in providing enhanced lubricity, mention may be made of such materials as copolymers of acrylic acid and a polyallyl sucrose, reaction products of cellulose or glucose with acids or alkylene oxides, natural gums, other synthetic gums, resins, latices, starches, alcohols, and protein gel-formers generally used as thickeners in cosmetic and pharmaceutical preparations. Particularly preferred water-soluble polymers include polyethylene oxide and hydroxyethyl cellulose (the latter sold as various grades of "Natrosol" by Hercules, Inc., Wilmington, Delaware). When included, the water-soluble polymer usually comprises 0.01 - 5%, preferably 0.1 - 2% by weight of the composition.

[0015] Other compatible additives as are well known in the art for use in shaving preparations may also be included in minor proportions so long as they do not adversely affect the properties of the gel. As examples of such additives, mention may be made of humectants, e.g., glycerine, sorbitol, glycereth -26; emollients, e.g., PEG-150 distearate, coco-triglycerides; skin freshening and skin soothing ingredients, e.g. menthol; aloe; lanolin; stability enhancers, e.g., lauramide DEA; perfume; colorants; opacifiers and clarifying agents as desired; and antiseptic agents.

[0016] The following Examples illustrate representative self-foaming shaving gel products and are given by way of

illustration only and are not to be considered as being limiting. The amounts in the Examples and the claims are in weight percent.

Example 1

[0017]

CTFA Name	% Active
Water	69.03
Palmitic Acid	8.17
Oleth-20	5.77
Triethanolamine (99%)	5.00
Isopentane	2.89
Isobutane	0.96
Hydrogenated Polyisobutenes (Panalane)	1.92
Glycereth-26	1.92
Lauramide DEA	1.44
Aloe Vera Gel	0.96
Fragrance	0.96
PEG-150 [®] Distearate	0.48
PEG-14M [®] (polyethylene oxide)	0.29
Hydroxyethyl Cellulose (Natrosol 250 HHR)	0.19
Acetic Acid (vehicle for fluorosurfactant)	0.005
Fluorosurfactant (Zonyl FSK)	0.005
Colorant	0.0006

[0018] Procedure: Form the aqueous phase by dissolving into the water at room temperature the Glycereth-26, hydroxyethyl cellulose, and PEG-14M[®]. Heat to 80° - 85°C., and add the palmitic acid, then triethanolamine with mixing. Form an oil, phase by mixing, at 55°C., the Oleth-20[®], PEG-150[®] distearate, hydrogenated polyisobutene, and lauramide DEA. Add the oil phase with thorough mixing to the aqueous phase maintained at 80°C. Cool to 40°C., and add the fluorosurfactant, aloe gel, fragrance, and dye. Mix and cool to room temperature; then blend with the mixed isopentane/isobutane and package in a barrier-type aerosol container.

[0019] The self-foaming shaving gel of Example 1 was tested against a commercial self-foaming gel well established in the marketplace, in a standard "crossover" test in which test panelists shave full face with one product for one week and then for another week with the other product. The identity of the test products is concealed from the test panelists. In this test the product of Example 1 was significantly preferred (at the 95% confidence level or better) in the key attributes of beard softening, avoidance of nicks and cuts, razor glide, and perception of facial cleanliness after shaving.

Example 2

[0020]

CTFA Name	% Active
Water	73.82

(continued)


CTFA Name	% Active
Stearic Acid	7.69
Oleth-20	5.76
Triethanolamine (99%)	3.37
Lauramide DEA	2.88
Isopentane	2.88
Isobutane	0.96
Hydrogenated Polyisobutene (Panalane®)	1.42
Fragrance	0.72
Hydroxyethyl Cellulose (Natrosol 250 HR®)	0.39
Coco-triglyceride	0.10
Fluorosurfactant (Zonyl FSK®)	0.005
Acetic Acid (vehicle for fluorosurfactant)	0.005
Colorant	0.0005

[0021] The procedure follows the process described in Example 1.

Claims

1. A shaving composition in the form of a self-foaming gel comprising water, a water-soluble soap, a volatile self-foaming agent, from 0.0005 - 0.5 percent by weight of a fluorosurfactant, and a hydrogenated polyisobutene.
2. The composition of claim 1, wherein the hydrogenated polyisobutene comprises from 0.5 - 5 percent by weight of the composition.
3. The composition of claim 2, wherein the fluoro-surfactant comprises from 0.001 - 0.1 percent by weight of the composition.
4. The composition of claim 2, wherein the fluoro-surfactant comprises from 0.002 - 0.01 percent by weight of the composition.
5. The composition of claim 2, which further comprises from 0.01 - 5 percent by weight of a water-soluble polymer.
6. The composition of claim 5, wherein the water-soluble polymer comprises from 0.1-2 percent by weight of the composition.
7. The composition of claim 6, wherein water comprises from 55 - 85 percent by weight, the soap comprises from 5 - 30 percent by weight, and the volatile self-foaming agent comprises from 0.5 - 10 percent by weight of the composition.
8. The composition of claim 5, wherein water comprises 65 - 80% in percent by weight water-soluble soap comprises 12-25% in percent by weight, water-soluble polymer comprises 0.1 - 2% in percent by weight, volatile self-foaming agent comprises 1 - 5%, fluorosurfactant comprises 0.002 - 0.01% in percent by weight, and hydrogenated polyisobutene comprises 1 - 4% in percent by weight.
9. The composition of claim 1, comprising in percent by weight of the composition, 65 - 80% water, 12 - 25% of a water soluble soap selected from sodium, potassium and triethanolamine salts of stearic, palmitic, myristic, oleic and coconut oil fatty acids, and mixtures thereof, 1 - 5% of a volatile self-foaming agent selected from n-pentane, isopentane, neopentane, n-butane, isobutane, and mixtures thereof, 0.002 - 0.01% of a fluorosurfactant, preferably not a cationic fluorosurfactant, and 1 - 4% of a hydrogenated polyisobutene.

10. The shaving composition of claim 9, wherein the fluorosurfactant is selected from



11. The shaving composition of claim 9, wherein the fluorosurfactant is $R_fCH_2CH(OCOCH_3)CH_2N^+(CH_3)_2CH_2CO_2^-$ wherein R_f is $F(CF_2CF_2)_{3-8}$.

12. The shaving composition of claim 1, additionally comprising a water soluble polyethoxylated fatty alcohol.

Patentansprüche

1. Zusammensetzung zum Rasieren in Form eines selbstschäumenden Gels, umfassend Wasser, eine wasserlösliche Seife, ein flüchtiges, selbstschäumendes Mittel, 0,0005 % ... 0,5 Gewichtsprozent eines Fluortensids und ein hydriertes Polyisobuten.

2. Zusammensetzung nach Anspruch 1, wobei das hydrierte Polyisobuten 0,5 % ... 5 Gewichtsprozent der Zusammensetzung ausmacht.

3. Zusammensetzung nach Anspruch 2, wobei das Fluortensid 0,001 % ... 0,1 Gewichtsprozent der Zusammensetzung ausmacht.

4. Zusammensetzung nach Anspruch 2, wobei das Fluortensid 0,002 % ... 0,01 Gewichtsprozent der Zusammensetzung ausmacht.

5. Zusammensetzung nach Anspruch 2, die ferner 0,01 % ... 5 Gewichtsprozent eines wasserlöslichen Polymers aufweist.


6. Zusammensetzung nach Anspruch 5, wobei das wasserlösliche Polymer 0,1 % ... 2 Gewichtsprozent der Zusammensetzung ausmacht.

7. Zusammensetzung nach Anspruch 6, wobei das Wasser 55 % ... 85 Gewichtsprozent ausmacht, die Seife 5 % ... 30 Gewichtsprozent ausmacht und das flüchtige, selbstschäumende Mittel 0,5 % ... 10 Gewichtsprozent der Zusammensetzung ausmacht.

8. Zusammensetzung nach Anspruch 5, wobei das Wasser 65 % ... 80 Gewichtsprozent ausmacht, wasserlösliche Seife 12 % ... 25 Gewichtsprozent ausmacht, wasserlösliches Polymer 0,1 % ... 2 Gewichtsprozent ausmacht, flüchtiges, selbstschäumendes Mittel 1 % ... 5 Gewichtsprozent ausmacht, Fluortensid 0,002 % ... 0,01, Gewichtsprozent ausmacht und hydriertes Polyisobuten 1 % ... 4 Gewichtsprozent ausmacht.

9. Zusammensetzung nach Anspruch 1, umfassend in Gewichtsprozent der Zusammensetzung: 65 ... 80 % Wasser; 12 ... 25 % wasserlösliche Seife, ausgewählt aus Natrium-, Kalium- und Triethanolamin-Salzen der Stearinsäure, Palmitinsäure, Myristinsäure, Öl- und Kokosölsäuren sowie Mischungen davon; 1 ... 5 % flüchtiges, selbstschäumendes Mittel, ausgewählt aus n-Pentan, Isopentan, Neopentan, n-Butan, Isobutan und Mischungen davon; 0,002 ... 0,01 % eines Fluortensids, vorzugsweise nicht ein kationisches Fluortensid; und 1 ... 4 % eines hydrierten Polyisobutens.

10. Zusammensetzung zum Rasieren nach Anspruch 9, wobei das Fluortensid ausgewählt wird aus:



11. Zusammensetzung zum Rasieren nach Anspruch 9, wobei das Fluortensid $R_fCH_2CH(OCOCH_3)CH_2N^+(CH_3)_2CH_2COO^-$ ist, worin R_f $F(CF_2CF_2)_{3...8}$ ist.

12. Zusammensetzung zum Rasieren nach Anspruch 1, die zusätzlich einen wasserlöslichen, polyethoxylierten Fettalkohol aufweist.

Revendications

5

1. Composition de rasage sous forme d'un gel automoussant comprenant de l'eau, un savon hydrosoluble, un agent automoussant volatil, 0,0005-0,5 % en masse d'un tensioactif fluoré et un polyisobutène hydrogéné.
2. Composition selon la revendication 1, dans laquelle le polyisobutène hydrogéné constitue 0,5-5 % en masse de la composition.
3. Composition selon la revendication 2, dans laquelle le tensioactif fluoré constitue 0,001-0,1 % en masse de la composition.
4. Composition selon la revendication 2, dans laquelle le tensioactif fluoré constitue 0,002-0,01 % en masse de la composition.
5. Composition selon la revendication 2 qui comprend en outre 0,01-5 % en masse d'un polymère hydrosoluble.
6. Composition selon la revendication 5, dans laquelle le polymère hydrosoluble constitue 0,1-2 % en masse de la composition.
7. Composition selon la revendication 6, dans laquelle l'eau constitue 55-85 % en masse, le savon constitue 5-30 % en masse et l'agent automoussant volatil constitue 0,5-10 % en masse de la composition.
8. Composition selon la revendication 5, dans laquelle l'eau constitue 65-80 % en masse, le savon hydrosoluble constitue 12-25 % en masse, le polymère hydrosoluble constitue 0,1-2 % en masse, l'agent automoussant volatil constitue 1-5 %, le tensioactif fluoré constitue 0,002-0,01 % en masse et le polyisobutène hydrogéné constitue 1-4 % en masse.
9. Composition selon la revendication 1 comprenant, en pourcentage en masse de la composition, 65-80 % d'eau, 12-25 % d'un savon hydrosoluble choisi parmi les sels de sodium, de potassium et de triéthanolamine d'acides gras stéarique, palmitique, myristique, oléique et d'huile de coprah, et leurs mélanges, 1-5 % d'un agent automoussant volatil choisi parmi le n-pentane, l'isopentane, le néopentane, le n-butane, l'isobutane, et leurs mélanges, 0,002-0,01 % d'un tensioactif fluoré, de préférence qui n'est pas un tensioactif fluoré cationique, et 1-4 % d'un polyisobutène hydrogéné.
10. Composition de rasage selon la revendication 9, dans laquelle le tensioactif fluoré est choisi parmi
 $R_fCH_2CH_2SCH_2CH_2CO_2Li$,
 $R_fCH_2CH(OCOCH_3)CH_2N^+(CH_3)_2CH_2CO_2^-$, et
 $R_fCH_2CH_2O(CH_2CH_2O)_xH$ où R_f est $F(CF_2CF_2)_{3-8}$.
11. Composition de rasage selon la revendication 9, dans laquelle le tensioactif fluoré est $R_fCH_2CH(OCOCH_3)CH_2N^+(CH_3)_2CH_2CO_2^-$ où R_f est $F(CF_2CF_2)_{3-8}$.
12. Composition de rasage selon la revendication 1, comprenant en outre un alcool gras polyéthoxylé hydrosoluble.

50

55

THIS PAGE BLANK (USPTO)